


Circle

Exercise 11A

Question 1:

Let AB be a chord of the given circle with centre O and radius 10 cm. Then, OA = 10 cm and AB = 16 cm. From O, draw OL ⊥ AB. We know that the perpendicular from the centre of a circle to a chord bisects the chord.

$$\begin{aligned}\therefore AL &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 16\right) \text{ cm} = 8 \text{ cm.}\end{aligned}$$



From right angled $\triangle OLA$, we have

$$\begin{aligned}OA^2 &= OL^2 + AL^2 \\ \Rightarrow OL^2 &= OA^2 - AL^2 \\ &= 10^2 - 8^2 \\ &= 100 - 64 = 36 \\ \therefore OL &= \sqrt{36} = 6 \text{ cm.}\end{aligned}$$

\therefore The distance of the chord from the centre is 6 cm.


Question 2:

Let AB be the chord of the given circle with centre O and radius 5 cm.

From O, draw OL ⊥ AB

Then, OA = 5 cm and OL = 3 cm [given]


We know that the perpendicular from the centre of a circle to a chord bisects the chord.



Now, in right angled $\triangle OLA$, we have

$$\begin{aligned}OA^2 &= AL^2 + OL^2 \\ \Rightarrow AL^2 &= OA^2 - OL^2 \\ \Rightarrow AL^2 &= 5^2 - 3^2 \\ &= 25 - 9 = 16 \\ \therefore AL &= \sqrt{16} = 4 \text{ cm} \\ \text{So, } AB &= 2 AL \\ &= (2 \times 4) \text{ cm} = 8 \text{ cm} \\ \therefore \text{ the length of the chord is 8 cm.}\end{aligned}$$

Let AB be the chord of the given circle with centre O. Draw $OL \perp AB$.



Then, OL is the distance from the centre to the chord.
So, we have $AB = 30$ cm and $OL = 8$ cm.

We know that the perpendicular from the centre of a circle to a chord bisects the chord.

$$\begin{aligned} \therefore AL &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 30\right) \text{ cm} = 15 \text{ cm} \end{aligned}$$

Now, in right angled $\triangle OLA$ we have,


$$\begin{aligned} OA^2 &= OL^2 + AL^2 \\ &= 8^2 + 15^2 \\ &= 64 + 225 = 289 \end{aligned}$$

$$\therefore OA = \sqrt{289} = 17 \text{ cm}$$

\therefore the radius of the circle is 17 cm.

Question 4:

(i) Let AB and CD be two chords of a circle such that $AB \parallel CD$ which are on the same side of the circle. Also $AB = 8$ cm and $CD = 6$ cm. $OB = OD = 5$ cm. Join OL and OM.
Since the perpendicular from the centre of a circle to a chord bisects the chord.



$$\begin{aligned} \text{We have } LB &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 8\right) \text{ cm} = 4 \text{ cm} \end{aligned}$$

$$\begin{aligned} \text{and } MD &= \frac{1}{2} \times CD \\ &= \left(\frac{1}{2} \times 6\right) \text{ cm} = 3 \text{ cm} \end{aligned}$$

Now in right angled $\triangle BLO$

$$\begin{aligned} OB^2 &= LB^2 + LO^2 \\ \Rightarrow LO^2 &= OB^2 - LB^2 \\ \Rightarrow &= 5^2 - 4^2 \\ &= 25 - 16 = 9 \end{aligned}$$

$$\therefore LO = \sqrt{9} = 3 \text{ cm.}$$


Again in right angled $\triangle DMO$

$$\begin{aligned} OD^2 &= MD^2 + MO^2 \\ \Rightarrow MO^2 &= OD^2 - MD^2 \\ &= 5^2 - 3^2 \\ &= 25 - 9 = 16 \end{aligned}$$

$$\Rightarrow MO = \sqrt{16} = 4 \text{ cm}$$

\therefore The distance between the chords $= (4 - 3) \text{ cm} = 1 \text{ cm}$.

- (ii) Let AB and CD be two chords of a circle such that $AB \parallel CD$ and they are on the opposite sides of the centre. $AB = 8$ cm and $CD = 6$ cm. Draw $OL \perp AB$ and $OM \perp CD$.



Join OA and OC

Then $OA = OC = 5$ cm (radius)

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have,

$$\begin{aligned} AL &= \frac{1}{2} AB \\ &= \left(\frac{1}{2} \times 8 \right) \text{ cm} = 4 \text{ cm.} \end{aligned}$$

$$\begin{aligned} \text{Also, } \quad CM &= \frac{1}{2} CD \\ &= \left(\frac{1}{2} \times 6 \right) \text{ cm} = 3 \text{ cm} \end{aligned}$$

Now, in right angled $\triangle OLA$, we have

$$\begin{aligned} OA^2 &= AL^2 + OL^2 \\ \Rightarrow \quad OL^2 &= OA^2 - AL^2 \\ &= 5^2 - 4^2 \\ &= 25 - 16 = 9 \text{ cm} \end{aligned}$$

$\therefore OL = \sqrt{9} = 3$ cm

Again in right angled $\triangle OMC$, we have


$$\begin{aligned} OC^2 &= OM^2 + CM^2 \\ \Rightarrow \quad OM^2 &= OC^2 - CM^2 \\ &= 5^2 - 3^2 \\ &= 25 - 9 = 16 \end{aligned}$$

$\Rightarrow OM = \sqrt{16} = 4$ cm

\therefore the distance between the chords $= (4 + 3)$ cm $= 7$ cm

Question 5:

Let AB and CD be two chords of a circle having centre O.
 AB = 30 cm and CD = 16 cm.



Join AO and OC which are its radii. So $AO = 17$ cm and $CO = 17$ cm.

Draw $OM \perp CD$ and $OL \perp AB$.

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have

$$\begin{aligned} AL &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 30 \right) \text{ cm} = 15 \text{ cm} \\ CM &= \frac{1}{2} \times CD \\ &= \left(\frac{1}{2} \times 16 \right) \text{ cm} = 8 \text{ cm} \end{aligned}$$

Now, in right-angled ΔALO , we have

$$\begin{aligned} AC^2 &= OL^2 + AL^2 \\ \Rightarrow LO^2 &= AC^2 - AL^2 \\ &= 17^2 - 15^2 \\ &= 289 - 225 = 64 \end{aligned}$$

$$\Rightarrow LO = \sqrt{64} = 8 \text{ cm}$$

Again, in right-angled ΔCMQ , we have


$$\begin{aligned} CO^2 &= CM^2 + OM^2 \\ \Rightarrow OM^2 &= CO^2 - CM^2 \\ &= 17^2 - 8^2 \\ &= 289 - 64 = 225 \end{aligned}$$

$$\Rightarrow OM = \sqrt{225} = 15 \text{ cm}$$

$$\begin{aligned} \therefore \text{Distance between the chords} &= OM + OL = (8 + 15) \text{ cm} \\ &= 23 \text{ cm.} \end{aligned}$$

Question 6:

CD is the diameter of a circle with centre O, and is perpendicular to chord AB. Join OA.



AB = 12 cm and CE = 3 cm [Given]

Let OA = OC = r cm

Then, OE = (r - 3) cm

Since the perpendicular from the centre of a circle to a chord bisects the chord, we have

$$\begin{aligned} AE &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 12\right) \text{ cm} = 6 \text{ cm} \end{aligned}$$

Now, in right angled $\triangle OEA$,

$$\begin{aligned} OA^2 &= OE^2 + AE^2 \\ \Rightarrow r^2 &= (r - 3)^2 + 6^2 \\ \Rightarrow &= r^2 - 6r + 9 + 36 \\ \Rightarrow r^2 - r^2 + 6r &= 45 \\ \Rightarrow 6r &= 45 \\ \Rightarrow r &= \frac{45}{6} = 7.5 \text{ cm} \end{aligned}$$

\therefore OA, the radius of the circle is 7.5 cm.

Question 7:


AB is the diameter of a circle with centre O which bisects the chord CD at point E.

CE = ED = 8 cm and EB = 4 cm. Join OC.

Let OC = OB = r cm.

Then,

OE = (r - 4) cm




Now, in right angled $\triangle OEC$

$$\begin{aligned} OC^2 &= OE^2 + EC^2 \\ r^2 &= (r - 4)^2 + 8^2 \\ \Rightarrow r^2 &= r^2 - 8r + 16 + 64 \\ \Rightarrow r^2 &= r^2 - 8r + 80 \\ \Rightarrow r^2 - r^2 + 8r &= 80 \\ \Rightarrow 8r &= 80 \\ \Rightarrow r &= \frac{80}{8} = 10 \text{ cm} \end{aligned}$$

\therefore the radius of the circle is 10 cm.

Question 8:

Given: $OD \perp AB$ of a circle with centre O . BC is a diameter.
 To Prove: $AC \parallel OD$ and $AC = 2 \times OD$
 Construction: Join AC .



Proof: We know that the perpendicular from the centre of the circle to a chord bisects the chord.

Here $OD \perp AB$

$\Rightarrow D$ is the mid-point of AB

$\Rightarrow AD = BD$

Also, O is the mid-point of BC

$\therefore OC = OB$

Now, in $\triangle ABC$, D is the midpoint of AB and O is the midpoint of BC

Midpoint Theorem: The line segment joining the midpoints of any two sides of a triangle is parallel to the third side and equal to half of it.

$\therefore OD \parallel AC$ and $OD = \frac{1}{2} AC$


$\therefore AC = 2 \times OD$

Question 9:

Sol.9: Given: O is the centre in which chords AB and CD intersect at P such that PO bisects $\angle BPD$.

To Prove: $AB = CD$

Construction: Draw $OE \perp AB$ and $OF \perp CD$



Proof: In $\triangle OEP$ and $\triangle OFP$

$\angle OEP = \angle OFP$ [Each equal to 90°]

$OP = OP$ [common]

$\angle OPE = \angle OPF$ [Since OP bisects $\angle BPD$]

Thus, by Angle-Side-Angle criterion of congruence, have,

$\therefore \triangle OEP \cong \triangle OFP$ [By ASA]

The corresponding parts of the congruent triangles are equal

$\Rightarrow OE = OF$ [C.P.C.T.]

\Rightarrow Chords AB and CD are equidistant from the centre O .


$\Rightarrow AB = CD$ [\because chords equidistant from the centre are equal]

$\therefore AB = CD$

Question 10:

Given: AB and CD are two parallel chords of a circle with centre O . POQ is a diameter which is perpendicular to AB .

To Prove: $PF \perp CD$ and $CF = FD$



Proof: $AB \parallel CD$ and POQ is a diameter.

$\angle PEB = 90^\circ$ [Given]

Then, $\angle PFD = \angle PEB$ [$AB \parallel CD$, Corresponding angles]

Thus, $PF \perp CD$

So, $OF \perp CD$


We know that, the perpendicular from the centre of a circle to chord, bisects the chord

$\therefore CF = FD$

Question 11:

If possible let two different circles intersect at three distinct point A, B and C.

Then, these points are noncollinear. So a unique circle can be drawn to pass through these points. This is a contradiction.

Question 12:

$$OA = 10 \text{ cm} \quad \text{and} \quad AB = 12 \text{ cm}$$

$$\therefore AD = \frac{1}{2} \times AB$$

$$AD = \left(\frac{1}{2} \times 12\right) \text{ cm} = 6 \text{ cm}$$

Now in right angled $\triangle ADO$,

$$OA^2 = AD^2 + OD^2$$

$$\Rightarrow OD^2 = OA^2 - AD^2$$

$$= 10^2 - 6^2$$

$$= 100 - 36 = 64$$

$$\therefore OD = \sqrt{64} = 8 \text{ cm}$$

Again, we have $O'A = 8 \text{ cm}$

In right angle $\triangle ADO'$

$$O'A^2 = AD^2 + O'D^2$$

$$\Rightarrow O'D^2 = O'A^2 - AD^2$$

$$= 8^2 - 6^2$$

$$= 64 - 36 = 28$$

$$O'D = \sqrt{28} = 2\sqrt{7} \text{ cm}$$

$$\therefore OO' = (OD + O'D)$$

$$= (8 + 2\sqrt{7}) \text{ cm}$$


\therefore the distance between their centres is $(8 + 2\sqrt{7}) \text{ cm}$.

Question 13:

Given: Two equal circles intersect at points P and Q. A straight line through P meets the circles in A and B.

To Prove: $QA = QB$

Construction: Join PQ.



Proof: Two circles will be congruent if and only if they have equal radii.

If two chords of a circle are equal then their corresponding arcs are congruent.

Here PQ is the common chord to both the circles.

Thus, their corresponding arcs are equal:


So, arc PCQ = arc PDQ

$\therefore \angle QAP = \angle QBP$ [congruent arcs have the same degree measure]

$\therefore QA = QB$ [isosceles triangle, base angles are equal]

Question 14:

Given: AB and CD are the two chords of a circle with centre O.
 Diameter POQ bisects them at L and M.
 To Prove: AB \parallel CD.



Proof: AB and CD are two chords of a circle with centre O.
 Diameter POQ bisects them at L and M.

Then, $OL \perp AB$
 and, $OM \perp CD$
 $\therefore \angle ALM = \angle LMD$
 $\therefore AB \parallel CD$ [alternate angles are equal]

Question 15:

Two circles with centres A and B, having radii 5 cm and 3 cm touch each other internally.

The perpendicular bisector of AB meets the bigger circle in P and Q.

Join AP.


Let PQ intersect AB at L.

Then, $AB = (5 - 3) \text{ cm} = 2 \text{ cm}$.

Since PQ is the perpendicular bisector of AB, we have

$$AL = \frac{1}{2} \times AB$$

$$= \left(\frac{1}{2} \times 2\right) \text{ cm} = 1 \text{ cm}$$



Now, in right angle $\triangle PLA$

$$\therefore AP^2 = AL^2 + PL^2$$

$$\Rightarrow PL = \sqrt{AP^2 - AL^2} \text{ cm}$$


$$= \sqrt{(25 - 1)} \text{ cm} = \sqrt{24} \text{ cm} = 2\sqrt{6} \text{ cm}$$

$$\therefore PQ = (2 \times PL) = (2 \times 2\sqrt{6}) \text{ cm} = 4\sqrt{6} \text{ cm}$$

$$\therefore \text{the length of } PQ = 4\sqrt{6} \text{ cm}$$

Question 16:


Given: AB is a chord of a circle with centre O. AB is produced to C such that BC = OB. Also, CO is joined to meet the circle in D. $\angle ACD = y^\circ$ and $\angle AOD = x^\circ$.



To Prove: $x = 3y$
 Proof: $OB = BC$ [Given]
 $\therefore \angle BOC = \angle BCO = y^\circ$ [Isosceles triangle]
 Ext. $\angle OBA = \angle BOC + \angle BCO = (2y)^\circ$
 Again, $OA = OB$ [radii of same circle]
 $\therefore \angle OAB = \angle OBA = (2y)^\circ$ [Isosceles triangle]
 Ext. $\angle AOD = \angle OAC + \angle ACO$
 $= \angle OAB + \angle BCO = 3y^\circ$
 $\therefore x^\circ = 3y^\circ$ [$\because \angle AOD = x$ (given)]

Question 17:

Given: AB and AC are chords of the circle with centre O. $AB = AC$, $OP \perp AB$ and $OQ \perp AC$.



To Prove: $PB = QC$
 Proof: $AB = AC$ [Given]
 $\therefore \frac{1}{2}AB = \frac{1}{2}AC$ [Divide by 2]

The perpendicular from the centre of a circle to a chord bisects the chord.

$\Rightarrow MB = NC \dots (1)$

Equal chords of a circle are equidistant from the centre.

$\Rightarrow OM = ON$

Also, $OP = OQ$ [Radii]

$\Rightarrow OP - OM = OQ - ON$

$\Rightarrow PM = QN \dots (2)$

Now consider the triangles, $\triangle MPB$ and $\triangle NQC$:

$MB = NC$ [from (1)]

$\angle PMB = \angle QNC$ [right angle, given]

$PM = QN$ [from (2)]

Thus, by Side-Angle-Side criterion of congruence, we have

$\therefore \triangle MPB \cong \triangle NQC$ [S.A.S.]

The corresponding parts of the congruent triangles are equal.

$\therefore PB = QC$ [by c.p.c.t.]

Question 18:

Given: BC is a diameter of a circle with centre O. AB and CD are two chords such that $AB \parallel CD$.

To Prove: $AB = CD$

Construction: Draw $OL \perp AB$ and $OM \perp CD$.

Proof: In $\triangle OLB$ and $\triangle OMC$

$$\angle OLB = \angle OMC \quad [\text{Perpendicular bisector, angle} = 90^\circ]$$

$$\angle OBL = \angle OCD \quad [AB \parallel CD, BC \text{ is a transversal, thus alternate interior angles are equal}]$$


$$OB = OC \quad [\text{Radii}]$$

Thus by Angle-Angle-Side criterion of congruence, we have

$$\therefore \triangle OLB \cong \triangle OMC \quad [\text{By AAS}]$$

The corresponding parts of the congruent triangle are equal.

$$\therefore OL = OM \quad [\text{C.P.C.T.}]$$



But the chords equidistant from the centre are equal.

$$\therefore AB = CD$$

Question 19:


Let $\triangle ABC$ be an equilateral triangle of side 9 cm.

Let AD be one of its medians.

Then, $AD \perp BC$.

$$\text{and } BD = \frac{1}{2} \times BC$$

$$= \left(\frac{1}{2} \times 9 \right) \text{ cm} = 4.5 \text{ cm.}$$



\therefore In right angled $\triangle ADB$,

$$AB^2 = AD^2 + BD^2$$

$$\Rightarrow AD^2 = AB^2 - BD^2$$

$$\Rightarrow AD = \sqrt{AB^2 - BD^2}$$

$$= \sqrt{(9)^2 - \left(\frac{9}{2}\right)^2} \text{ cm} = \frac{9\sqrt{3}}{2} \text{ cm}$$

In an equilateral triangle, the centroid and circumcentre coincide and $AG : GD = 2 : 1$

$$\therefore \text{radius } AG = \frac{2}{3} AD$$

$$= \left(\frac{2}{3} \times \frac{9\sqrt{3}}{2} \right) \text{ cm} = 3\sqrt{3} \text{ cm}$$


\therefore The radius of the circle is $3\sqrt{3}$ cm.

Question 20:

Given : AB and AC are two equal chords of a circle with centre O

To Prove: $\angle OAB = \angle OAC$

Construction: Join OA, OB and OC.



Proof: In $\triangle OAB$ and $\triangle OAC$,

$$AB = AC \quad \text{[Given]}$$

$$OA = OA \quad \text{[common]}$$

$$OB = OC \quad \text{[Radii]}$$

Thus by Side-Side-Side criterion of congruence, we have

$$\therefore \triangle OAB \cong \triangle OAC \quad \text{[by SSS]}$$

The corresponding parts of the congruent triangles are equal.


$$\Rightarrow \angle OAB = \angle OAC \quad \text{[by C.P.C.T.]}$$

Therefore, O lies on the bisector of $\angle BAC$.

Question 21:

Given: OPQR is a square. A circle with centre O cuts the square in X and Y.

To Prove: $QX = QY$



Construction: Join OX and OY.

Proof: In $\triangle OXP$ and $\triangle OYR$,

$$\angle OPX = \angle ORY \quad \text{[Each equal to } 90^\circ \text{]}$$

$$OX = OY \quad \text{[Radii]}$$

$$OP = OR \quad \text{[Sides of a square]}$$

Thus by Right Angle-Hypotenuse-Side criterion of congruence, we have,

$$\therefore \triangle OXP \cong \triangle OYR \quad \text{[by RHS]}$$

The corresponding parts of the congruent triangles are equal.

$$\Rightarrow PX = RY \quad \text{[by C.P.C.T.]}$$

$$\Rightarrow PQ - PX = QR - RY \quad \text{[} \because PQ = QR \text{]}$$

$$\therefore QX = QY.$$

Exercise 11B

Question 1:

(i) Join BO.

In $\triangle BOC$ we have


$$OC = OB \quad \left[\begin{array}{l} \text{Each equal to the radius} \end{array} \right]$$

$$\Rightarrow \angle OBC = \angle OCB \quad \left[\begin{array}{l} \text{Base angles of an isosceles} \\ \text{triangle are equal} \end{array} \right]$$

$$\Rightarrow \angle OBC = 30^\circ \quad \left[\begin{array}{l} \angle OCB = 30^\circ \end{array} \right]$$

Thus, we have,

$$\angle OBC = 30^\circ \quad \dots\dots(1)$$



Now, in $\triangle BOA$, we have

$$OB = OC \quad [\text{Each equal to the radius}]$$

$$\Rightarrow \angle OAB = \angle OBA \quad [\text{base angles of an isosceles triangle are equal}]$$

$$\Rightarrow \angle OBA = 40^\circ \quad [\angle OAB = 40^\circ, \text{ given}]$$

Thus, we have,

$$\angle OBA = 40^\circ \quad \dots (2)$$

$$\therefore \angle ABC = \angle OBC + \angle OBA$$

$$\Rightarrow = 30^\circ + 40^\circ \quad [\text{from (1) and (2)}]$$


$$\Rightarrow \angle ABC = 70^\circ$$

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference,

$$\therefore \angle AOC = 2 \times \angle ABC \\ = 2 \times 70^\circ = 140^\circ$$

$$(ii) \quad \angle BOC = 360^\circ - (\angle AOB + \angle AOC) \\ = 360^\circ - (90^\circ + 110^\circ) \\ = 360^\circ - 200^\circ = 160^\circ$$

We know that $\angle BOC = 2\angle BAC$



$$\Rightarrow \angle BAC = \frac{160^\circ}{2} = 80^\circ \quad [\because \angle BOC = 160^\circ]$$

$$\therefore \angle BAC = 80^\circ$$


Question 2:

(i)

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\therefore \angle AOB = 2\angle OCA$$

$$\Rightarrow \angle OCA = \frac{70}{2} = 35^\circ \quad [\because \angle AOB = 70^\circ]$$




(ii) The radius of the circle is

$$OA = OC$$

$$\Rightarrow \angle OAC = \angle OCA \quad [\text{base angles of an isosceles triangle are equal}]$$

$$\Rightarrow \angle OAC = 35^\circ \quad [\text{as } \angle OCA = 35^\circ]$$

Question 3:



It is clear that $\angle ACB = \angle PCB$.

Consider the triangle $\triangle PCB$.

Applying the angle sum property, we have,

$$\angle PCB = 180^\circ - (\angle BPC + \angle PBC)$$

$$= 180^\circ - (180^\circ - 110^\circ + 25^\circ) \quad [\angle APB \text{ and } \angle BPC \text{ are linear pair; } \angle PBC = 25^\circ, \text{ given}]$$


$$= 180^\circ - (70^\circ + 25^\circ)$$

$$\angle PCB = 180^\circ - 95^\circ = 85^\circ$$

Angles in the same segment of a circle are equal.

$$\therefore \angle ADB = \angle ACB = 85^\circ$$

Question 4:



It is clear that, BD is the diameter of the circle.

Also we know that, the angle in a semicircle is a right angle.

$$\therefore \angle BAD = 90^\circ$$

Now consider the triangle, $\triangle BAD$

$$\Rightarrow \angle ADB = 180^\circ - (\angle BAD + \angle ABD) \quad [\text{Angle sum property}]$$

$$\Rightarrow \quad = 180^\circ - (90^\circ + 35^\circ) \quad [\angle BAD = 90^\circ \text{ and } \angle ABD = 35^\circ]$$

$$\Rightarrow \quad = 180^\circ - 125^\circ$$


$$\Rightarrow \angle ADB = 55^\circ$$

Angles in the same segment of a circle are equal.

$$\therefore \angle ACB = \angle ADB = 55^\circ$$

$$\therefore \angle ACB = 55^\circ$$

Question 5:



The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\begin{aligned} \therefore \angle AOB &= 2\angle ACB \\ &= 2 \times 50^\circ \quad [\text{Given}] \\ \Rightarrow \angle AOB &= 100^\circ \quad \dots(1) \end{aligned}$$

Consider the triangle $\triangle OAB$.

$$\begin{aligned} OA &= OB && [\text{radius of the circle}] \\ \angle OAB &= \angle OBA && [\text{base angles of an isosceles triangle are equal}] \end{aligned}$$

Thus we have

$$\angle OAB = \angle OBA \quad \dots(2)$$

By angle sum property, we have

$$\text{Now } \angle AOB + \angle OAB + \angle OBA = 180^\circ$$


$$\Rightarrow 100^\circ + 2\angle OAB = 180^\circ \quad [\text{from (1) and (2)}]$$

$$\Rightarrow 2\angle OAB = 180^\circ - 100^\circ = 80^\circ$$

$$\Rightarrow \angle OAB = \frac{80^\circ}{2} = 40^\circ$$

$$\therefore \angle OAB = 40^\circ$$

Question 6:



(i) Angles in the same segment of a circle are equal.
 $\angle ABD$ and $\angle ACD$ are in the segment AD.

$$\begin{aligned} \therefore \angle ACD &= \angle ABD \\ &= 54^\circ \quad [\text{Given}] \end{aligned}$$

(ii) Angles in the same segment of a circle are equal.
 $\angle BAD$ and $\angle BCD$ are in the segment BD.

$$\begin{aligned} \therefore \angle BAD &= \angle BCD \\ &= 43^\circ \quad [\text{Given}] \end{aligned}$$

(iii) Consider the $\triangle ABD$.

By Angle sum property we have


$$\angle BAD + \angle ADB + \angle DBA = 180^\circ$$

$$\Rightarrow 43^\circ + \angle ADB + 54^\circ = 180^\circ$$

$$\Rightarrow \angle ADB = 180^\circ - 97^\circ = 83^\circ$$

$$\Rightarrow \angle BDA = 83^\circ$$

Question 7:



Angles in the same segment of a circle are equal.
 $\angle CAD$ and $\angle CBD$ are in the segment CD.

$$\therefore \angle CAD = \angle CBD = 60^\circ \quad [\text{Given}]$$

We know that an angle in a semi circle is a right angle:

$$\therefore \angle ADC = 90^\circ \quad [\text{angle in a semicircle}]$$

$$\begin{aligned} \therefore \angle ACD &= 180^\circ - (\angle ADC + \angle CAD) \\ &= 180^\circ - (90^\circ + 60^\circ) \\ &= 180^\circ - 150^\circ = 30^\circ \end{aligned}$$

$$\therefore \angle CDE = \angle ACD = 30^\circ \quad \left[\begin{array}{l} AC \parallel DE \text{ and } CD \text{ is a transversal,} \\ \text{thus alternate angles are equal} \end{array} \right]$$

Question 8:



Join CO and DO, $\angle BCD = \angle ABC = 25^\circ$ [alternate interior angles]

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\begin{aligned} \therefore \angle BOD &= 2\angle BCD \\ &= 50^\circ \quad [\angle BCD = 25^\circ] \end{aligned}$$

Similarly,

$$\begin{aligned} \angle AOC &= 2\angle ABC \\ &= 50^\circ \end{aligned}$$

AB is a straight line passing through the centre.

$$\therefore \angle AOC + \angle COD + \angle BOD = 180^\circ$$

$$\Rightarrow 50^\circ + \angle COD + 50^\circ = 180^\circ$$

$$\Rightarrow \angle COD = 180^\circ - 100^\circ = 80^\circ$$

$$\begin{aligned} \therefore \angle CED &= \frac{1}{2} \angle COD \\ &= \frac{80^\circ}{2} = 40^\circ \end{aligned}$$

$$\therefore \angle CED = 40^\circ$$

Question 9:

$$(i) \angle CED = 90^\circ$$


In $\triangle CED$, we have

$$\angle CED + \angle EDC + \angle DCE = 180^\circ$$

$$\Rightarrow 90^\circ + 40^\circ + \angle DCE = 180^\circ$$

$$\therefore \angle DCE = 180^\circ - 130^\circ$$

$$\angle DCE = 50^\circ \quad \dots(1)$$



(ii) $\angle AOC$ and $\angle BOC$ are linear pair.


$$\therefore \angle BOC = (180^\circ - 80^\circ) = 100^\circ \quad \dots(2)$$

$$\therefore \angle ABC = 180^\circ - (\angle BOC + \angle DCE)$$

$$= 180^\circ - (100^\circ + 50^\circ) \quad [\text{from (1) and (2)}]$$

$$= 180^\circ - 150^\circ = 30^\circ$$

Question 10:



The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\begin{aligned} \therefore \angle AOB &= 2\angle ACB \\ \Rightarrow &= 2\angle DCB \quad [\because \angle ACB = \angle DCB] \end{aligned}$$

$$\begin{aligned} \Rightarrow \angle DCB &= \frac{1}{2} \angle AOB \\ &= \left(\frac{1}{2} \times 40\right) = 20^\circ \end{aligned}$$

Consider the $\triangle OBC$;

By angle sum property, we have

$$\begin{aligned} \angle BOC + \angle OCB + \angle OBC &= 180^\circ \\ \Rightarrow 100^\circ + 20^\circ + \angle OBC &= 180^\circ \\ \Rightarrow \angle OBC &= 180^\circ - 120^\circ = 60^\circ \\ \Rightarrow \angle OBC &= \angle OCB = 60^\circ \\ \therefore \angle OBC &= 60^\circ \end{aligned}$$

Question 11:

Join OB.


$$\begin{aligned} \therefore OA &= OB \quad [\text{Radius}] \\ \therefore \angle OBA &= \angle OAB = 25^\circ \quad [\text{base angles are equal in isosceles triangle}] \end{aligned}$$

Now in $\triangle OAB$, we have

$$\begin{aligned} \Rightarrow \angle OAB + \angle OBA + \angle AOB &= 180^\circ \\ \Rightarrow 25^\circ + 25^\circ + \angle AOB &= 180^\circ \\ \Rightarrow \angle AOB &= 180^\circ - 50^\circ = 130^\circ \end{aligned}$$

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\begin{aligned} \therefore \angle AOB &= 2\angle ACB \\ \Rightarrow \angle ACB &= \frac{1}{2} \angle AOB = \frac{1}{2} \times 130 = 65^\circ \\ \Rightarrow \angle ECB &= 65^\circ \end{aligned}$$



Consider the right triangle $\triangle BEC$.

We know that the sum of three angles in a triangle is 180° .

$$\begin{aligned} \Rightarrow \angle EBC + \angle BEC + \angle ECB &= 180^\circ \\ \Rightarrow \angle EBC + 90^\circ + 65^\circ &= 180^\circ \\ \Rightarrow \angle EBC &= 180^\circ - 155^\circ = 25^\circ \\ \therefore \angle EBC &= 25^\circ \end{aligned}$$

Question 12:


$$\begin{aligned} & \Rightarrow OB = OC \quad \text{[Radius]} \\ & \Rightarrow \angle OBC = \angle OCB = 55^\circ \quad \text{[base angles in an isosceles triangle are equal]} \end{aligned}$$

Consider the triangle $\triangle BOC$.

By angle sum property, we have

$$\begin{aligned} \angle BOC &= 180^\circ - (\angle OCB + \angle OBC) \\ &= 180^\circ - (55^\circ + 55^\circ) \\ &= 180^\circ - 110^\circ = 70^\circ \end{aligned}$$

$$\therefore \angle BOC = 70^\circ$$



$$\begin{aligned} \text{Again, } & OA = OB \\ \Rightarrow & \angle OBA = \angle OAB = 20^\circ \quad \text{[base angles in an isosceles triangle are equal]} \end{aligned}$$

Consider the triangle $\triangle AOB$.


By angle sum property, we have

$$\begin{aligned} \Rightarrow \angle AOB &= 180^\circ - (\angle OAB + \angle OBA) \\ &= 180^\circ - (20^\circ + 20^\circ) \\ &= 180^\circ - 40^\circ = 140^\circ \end{aligned}$$

$$\begin{aligned} \therefore \angle AOC &= \angle AOB - \angle BOC \\ &= 140^\circ - 70^\circ = 70^\circ \end{aligned}$$

$$\therefore \angle AOC = 70^\circ$$

Question 13:



Join OB and OC.

The angle subtended by an arc of a circle at the centre is double the angle subtended by the arc at any point on the circumference.

$$\begin{aligned} \therefore \angle BOC &= 2\angle BAC \\ &= 2 \times 30^\circ \quad \text{[} \angle BAC = 30^\circ \text{]} \\ &= 60^\circ \quad \dots\dots(1) \end{aligned}$$

Now consider the triangle $\triangle BOC$.

$$\begin{aligned} & OB = OC \quad \text{[radii]} \\ \Rightarrow & \angle OBC = \angle OCB \quad \dots\dots(2) \end{aligned} \quad \left. \begin{array}{l} \text{base angles in an isosceles triangle} \\ \text{are equal} \end{array} \right\}$$

Now, in $\triangle BOC$, we have

$$\begin{aligned} \angle BOC + \angle OBC + \angle OCB &= 180^\circ \\ \Rightarrow 60^\circ + \angle OCB + \angle OCB &= 180^\circ \quad \text{[from (1) and (2)]} \\ \Rightarrow 2\angle OCB &= 180^\circ - 60^\circ \\ \Rightarrow &= 120^\circ \\ \Rightarrow \angle OCB &= \frac{120^\circ}{2} = 60^\circ \\ \Rightarrow \angle OBC &= 60^\circ \quad \text{[from (2)]} \end{aligned}$$

Thus, we have, $\angle OBC = \angle OCB = \angle BOC = 60^\circ$

So, $\triangle BOC$ is an equilateral triangle

$$\Rightarrow OB = OC = BC$$

$\therefore BC$ is the radius of the circumference.

Question 14:

Consider the triangle, ΔPRQ ,

PQ is the diameter.

The angle in a semicircle is a right angle:

$$\Rightarrow \angle PRQ = 90^\circ$$

By the angle sum property in ΔPRQ , we have,

$$\angle QPR + \angle PRQ + \angle PQR = 180^\circ$$

$$\Rightarrow \angle QPR + 90^\circ + 65^\circ = 180^\circ$$

$$\Rightarrow \angle QPR = 180^\circ - 155^\circ = 25^\circ \dots\dots(1)$$



Now consider the triangle ΔPQM .

Since PQ is the diameter, $\angle PMQ = 90^\circ$

Again applying the angle sum property in ΔPQM , we have

$$\angle QPM + \angle PMQ + \angle PQM = 180^\circ$$

$$\Rightarrow \angle QPM + 90^\circ + 50^\circ = 180^\circ$$

$$\Rightarrow \angle QPM = 180^\circ - 140^\circ = 40^\circ$$

Now in quadrilateral PQRS

$$\angle QPS + \angle SRQ = 180^\circ$$

$$\Rightarrow \angle QPR + \angle RPS + \angle PRQ + \angle PRS = 180^\circ \quad [\text{from (1)}]$$

$$\Rightarrow 25^\circ + 40^\circ + 90^\circ + \angle PRS = 180^\circ$$

$$\Rightarrow \angle PRS = 180^\circ - 155^\circ = 25^\circ$$

$$\therefore \angle PRS = 25^\circ$$

Exercise 11C

Question 1:

$$\angle BDC = \angle BAC = 40^\circ \text{ [angles in the same segment]}$$


In $\triangle BCD$, we have

$$\angle BCD + \angle BDC + \angle DBC = 180^\circ$$

$$\therefore \angle BCD + 40^\circ + 60^\circ = 180^\circ$$

$$\Rightarrow \angle BCD = 180^\circ - 100^\circ = 80^\circ$$

$$\therefore \angle BCD = 80^\circ$$



$$(ii) \text{ Also } \angle CAD = \angle CBD \text{ [angles in the same segment]}$$


$$\therefore \angle CAD = 60^\circ \text{ [}\because \angle CBD = 60^\circ\text{]}$$

Question 2:

In cyclic quadrilateral PQRS

$$\begin{aligned} \angle PSR + \angle PQR &= 180^\circ \\ \Rightarrow 150^\circ + \angle PQR &= 180^\circ \\ \Rightarrow \angle PQR &= 180^\circ - 150^\circ = 30^\circ \dots\dots (i) \\ \text{Also, } \angle PRQ &= 90^\circ \dots\dots (ii) \end{aligned}$$

[angle in a semi circle]




Now in $\triangle PRQ$ we have

$$\begin{aligned} \angle PQR + \angle PRQ + \angle RPQ &= 180^\circ \\ \Rightarrow 30^\circ + 90^\circ + \angle RPQ &= 180^\circ \text{ [from (i) and (ii)]} \\ \Rightarrow \angle RPQ &= 180^\circ - 120^\circ = 60^\circ \\ \therefore \angle RPQ &= 60^\circ \end{aligned}$$

Question 3:

In cyclic quadrilateral ABCD, $AB \parallel DC$ and $\angle BAD = 100^\circ$



$$\begin{aligned} (i) \quad \angle BCD + \angle BAD &= 180^\circ \\ \Rightarrow \angle BCD + 100^\circ &= 180^\circ \\ \Rightarrow \angle BCD &= 180^\circ - 100^\circ = 80^\circ \\ (ii) \quad \text{Also, } \angle ADC &= \angle BCD = 80^\circ \\ \therefore \angle ADC &= 80^\circ \\ (iii) \quad \angle ABC &= \angle BAD = 100^\circ \\ \therefore \angle ABC &= 100^\circ \end{aligned}$$


Question 4:

Take a point D on the major arc CA and join AD and DC

$$\therefore \angle 2 = 2\angle 1$$

[Angle subtended by an arc is twice the angle subtended by it on the circumference in the alternate segment.]

$$\begin{aligned} \therefore 130^\circ &= 2\angle 1 \\ \Rightarrow \angle 1 &= 65^\circ \dots\dots (i) \end{aligned}$$



$$\angle PBC = \angle 1$$


[\therefore exterior angle of a cyclic quadrilateral interior opposite angle]

$$\therefore \angle PBC = 65^\circ$$

Question 5:

ABCD is a cyclic quadrilateral

$$\begin{aligned} \therefore \angle ABC + \angle ADC &= 180^\circ \\ \Rightarrow 92^\circ + \angle ADC &= 180^\circ \\ \Rightarrow \angle ADC &= 180^\circ - 92^\circ = 88^\circ \end{aligned}$$




Also, $AE \parallel CD$

$$\begin{aligned} \therefore \angle EAD &= \angle ADC = 88^\circ \\ \angle BCD &= \angle DAF \\ \text{[exterior angle of a cyclic quadrilateral = int. opp. angle]} \\ \therefore \angle BCD &= \angle EAD + \angle EAF \\ &= 88^\circ + 20^\circ \quad [\because \angle FAE = 20^\circ \text{ (given)}] \\ &= 108^\circ \\ \therefore \angle BCD &= 108^\circ \end{aligned}$$

Question 6:

$$\begin{aligned} BD &= DC \\ \therefore \angle BCD &= \angle CBD = 30^\circ \end{aligned}$$



In $\triangle BCD$, we have

$$\begin{aligned} \angle BCD + \angle CBD + \angle CDB &= 180^\circ \\ \Rightarrow 30^\circ + 30^\circ + \angle CDB &= 180^\circ \\ \Rightarrow \angle CDB &= 180^\circ - 60^\circ \\ &= 120^\circ \end{aligned}$$

The opposite angles of a cyclic quadrilateral are supplementary.
ABCD is a cyclic quadrilateral and thus,

$$\begin{aligned} \angle CDB + \angle BAC &= 180^\circ \\ &= 180^\circ - 120^\circ \quad [\because \angle CDB = 120^\circ] \\ &= 60^\circ \\ \therefore \angle BAC &= 60^\circ \end{aligned}$$

Question 7:


Angle subtended by an arc is twice the angle subtended by it on the circumference in the alternate segment.
 Here arc ABC makes $\angle AOC = 100^\circ$ at the centre of the circle and $\angle ADC$ on the circumference of the circle

$$\therefore \angle AOC = 2\angle ADC$$

$$\Rightarrow \angle ADC = \frac{1}{2}(\angle AOC)$$

$$\Rightarrow \quad = \frac{1}{2} \times 100^\circ \quad [\angle AOC = 100^\circ]$$

$$\Rightarrow \angle ADC = 50^\circ$$



The opposite angles of a cyclic quadrilateral are supplementary.
 ABCD is a cyclic quadrilateral and thus,

$$\begin{aligned} \angle ADC + \angle ABC &= 180^\circ \\ &= 180^\circ - 50^\circ \quad [\angle ADC = 50^\circ] \\ &= 130^\circ \end{aligned}$$

$$\therefore \angle ABC = 130^\circ$$


$$\therefore \angle ADC = 50^\circ \text{ and } \angle ABC = 130^\circ$$

Question 8:

$\triangle ABC$ is an equilateral triangle.

\therefore Each of its angle is equal to 60°

$$\Rightarrow \angle BAC = \angle ABC = \angle ACB = 60^\circ$$



(i) Angles in the same segment of a circle are equal.

$$\therefore \angle BDC = \angle BAC$$

$$= 60^\circ \quad [\angle BAC = 60^\circ]$$

$$\Rightarrow \angle BDC = 60^\circ$$

(ii) The opposite angles of a cyclic quadrilateral are supplementary.

ABCE is a cyclic quadrilateral and thus,

$$\angle BAC + \angle BEC = 180^\circ$$

$$\angle BEC = 180^\circ - 60^\circ \quad [\angle BAC = 60^\circ]$$

$$= 120^\circ$$

$$\Rightarrow \angle BEC = 120^\circ$$


Question 9:

ABCD is a cyclic quadrilateral.

$$\therefore \angle A + \angle C = 180^\circ \quad \left\{ \begin{array}{l} \text{opp. angle of a cyclic quadrilateral} \\ \text{are supplementary} \end{array} \right.$$

$$\Rightarrow \angle A + 100^\circ = 180^\circ$$

$$\Rightarrow \angle A = 180^\circ - 100^\circ = 80^\circ$$



Now in $\triangle ABD$, we have

$$\angle A + \angle ABD + \angle ADB = 180^\circ$$


$$\Rightarrow 80^\circ + 50^\circ + \angle ADB = 180^\circ$$

$$\Rightarrow \angle ADB = 180^\circ - 130^\circ = 50^\circ$$

$$\therefore \angle ADB = 50^\circ$$

Question 10:

O is the centre of the circle and $\angle BOD = 150^\circ$
 \therefore Reflex $\angle BOD = (360^\circ - \angle BOD)$
 $= (360^\circ - 150^\circ) = 210^\circ$




Now, $x = \frac{1}{2}(\text{reflex } \angle BOD)$
 $= \frac{1}{2} \times 210^\circ = 105^\circ$

$\therefore x = 105^\circ$
 Again, $x + y = 180^\circ$
 $\Rightarrow 105^\circ + y = 180^\circ$
 $\Rightarrow y = 180^\circ - 105^\circ = 75^\circ$
 $\therefore y = 75^\circ$

Question 11:

O is the centre of the circle and $\angle DAB = 50^\circ$
 $OA = OB$ [Radii]
 $\Rightarrow \angle OBA = \angle OAB = 50^\circ$




In $\triangle OAB$ we have
 $\angle OAB + \angle OBA + \angle AOB = 180^\circ$
 $\Rightarrow 50^\circ + 50^\circ + \angle AOB = 180^\circ$
 $\Rightarrow \angle AOB = 180^\circ - 100^\circ = 80^\circ$
 Since, AOD is a straight line,
 $\therefore x = 180^\circ - \angle AOB$
 $= 180^\circ - 80^\circ = 100^\circ$
 $\therefore x = 100^\circ$

The opposite angles of a cyclic quadrilateral are supplementary.
 ABCD is a cyclic quadrilateral and thus,
 $\angle DAB + \angle BCD = 180^\circ$
 $\angle BCD = 180^\circ - 50^\circ$ [$\because \angle DAB = 50^\circ$, given]
 $= 130^\circ$
 $\Rightarrow y = 130^\circ$
 Thus, $x = 100^\circ$ and $y = 130^\circ$

Question 12:

ABCD is a cyclic quadrilateral.
 We know that in a cyclic quadrilateral exterior angle = interior opposite angle.
 $\therefore \angle CBF = \angle CDA = (180^\circ - x)$
 $\Rightarrow 130^\circ = 180^\circ - x$
 $\Rightarrow x = 180^\circ - 130^\circ = 50^\circ$
 $x = 50^\circ$



Question 13:

AB is a diameter of a circle with centre O and $DO \parallel CB$.


$$\angle BCD = 120^\circ$$

(i) Since ABCD is a cyclic quadrilateral.

$$\therefore \angle BCD + \angle BAD = 180^\circ$$

$$\Rightarrow 120^\circ + \angle BAD = 180^\circ$$

$$\Rightarrow \angle BAD = 180^\circ - 120^\circ = 60^\circ$$



(ii) $\angle BDA = 90^\circ$ [angle in a semi circle]

In $\triangle ABD$ we have

$$\angle BDA + \angle BAD + \angle ABD = 180^\circ$$

$$\Rightarrow 90^\circ + 60^\circ + \angle ABD = 180^\circ$$

$$\Rightarrow \angle ABD = 180^\circ - 150^\circ = 30^\circ$$

(iii) $OD = OA$,

$$\Rightarrow \angle ODA = \angle OAD = \angle BAD = 60^\circ$$

$$\therefore \angle ODB = 90^\circ - \angle ODA$$

$$= 90^\circ - 60^\circ = 30^\circ$$

Since $DO \parallel CB$, alternate angles are equal

$$\Rightarrow \angle CBD = \angle ODB$$

$$= 30^\circ$$

(iv) $\angle ADC = \angle ADB + \angle CDB$

$$= 90^\circ + 30^\circ = 120^\circ$$

Also, in $\triangle AOD$, we have

$$\angle ODA + \angle OAD + \angle AOD = 180^\circ$$

$$\Rightarrow 60^\circ + 60^\circ + \angle AOD = 180^\circ$$

$$\Rightarrow \angle AOD = 180^\circ - 120^\circ = 60^\circ$$

Since all the angles of $\triangle AOD$ are of 60° each

$\therefore \triangle AOD$ is an equilateral triangle.


Question 14:

AB and CD are two chords of a circle which intersect each other at

P, outside the circle. $AB = 6$ cm, $BP = 2$ cm and $PD = 2.5$ cm

Therefore, $AP \times BP = CP \times DP$

$$\text{Or, } 8 \times 2 = (CD + 2.5) \times 2.5 \text{ cm [as } CP = CD + DP]$$



Let $x = CD$

$$\text{Thus, } 8 \times 2 = (x + 2.5) \times 2.5$$

$$\Rightarrow 16 \text{ cm} = 2.5x + 6.25 \text{ cm}$$

$$\Rightarrow 2.5x = (16 - 6.25) \text{ cm}$$

$$\Rightarrow 2.5x = 9.75 \text{ cm}$$

$$\Rightarrow x = \frac{9.75}{2.5} = 3.9 \text{ cm}$$

$$\therefore x = 3.9 \text{ cm}$$

Therefore, $CD = 3.9$ cm


Question 15:

O is the centre of a circle having $\angle AOD = 140^\circ$ and $\angle CAB = 50^\circ$

$$(i) \quad \begin{aligned} \angle BOD &= 180^\circ - \angle AOD \\ &= 180^\circ - 140^\circ = 40^\circ \end{aligned}$$

$$OB = OD$$

$$\therefore \angle OBD = \angle ODB$$



In $\triangle OBD$, we have

$$\angle BOD + \angle OBD + \angle ODB = 180^\circ$$

$$\Rightarrow \angle BOD + \angle OBD + \angle OBD = 180^\circ \quad [\because \angle OBD = \angle ODB]$$

$$\Rightarrow 40^\circ + 2\angle OBD = 180^\circ \quad [\because \angle BOD = 40^\circ]$$

$$\Rightarrow 2\angle OBD = 180^\circ - 40^\circ = 140^\circ$$

$$\Rightarrow \angle OBD = \angle ODB = \frac{140^\circ}{2} = 70^\circ$$

$$\text{Also, } \angle CAB + \angle BDC = 180^\circ \quad [\because ABCD \text{ is cyclic}]$$

$$\Rightarrow \angle CAB + \angle ODB + \angle ODC = 180^\circ$$

$$\Rightarrow 50^\circ + 70^\circ + \angle ODC = 180^\circ$$

$$\Rightarrow \angle ODC = 180^\circ - 120^\circ = 60^\circ$$

$$\therefore \angle ODC = 60^\circ$$

$$\therefore \angle EDB = 180^\circ - (\angle ODC + \angle ODB)$$

$$= 180^\circ - (60^\circ + 70^\circ)$$

$$= 180^\circ - 130^\circ = 50^\circ$$


$$(ii) \quad \begin{aligned} \angle EBD &= 180^\circ - \angle OBD \\ &= 180^\circ - 70^\circ = 110^\circ \end{aligned}$$

Question 16:

Consider the triangles, $\triangle EBC$ and $\triangle EDA$

Side AB of the cyclic quadrilateral ABCD is produced to E

$$\begin{aligned} \therefore \angle EBC &= \angle CDA \\ \Rightarrow \angle EBC &= \angle EDA \quad \dots (i) \end{aligned}$$



Again, side DC of the cyclic quadrilateral ABCD is produced to E,

$$\begin{aligned} \therefore \angle ECB &= \angle BAD \\ \Rightarrow \angle ECB &= \angle EAD \quad \dots (ii) \end{aligned}$$

$$\text{and } \angle BEC = \angle DEA \quad [\text{each equal to } \angle E] \dots (iii)$$

Thus from (i), (ii) and (iii), we have

$$\triangle EBC \cong \triangle EDA$$

Question 17:

$\triangle ABC$ is an isosceles triangle in which $AB = AC$ and a circle passing through B and C intersects AB and AC at D and E.


$$\text{Since } AB = AC$$

$$\therefore \angle ACB = \angle ABC$$

$$\text{So, ext. } \angle ADE = \angle ACB = \angle ABC$$


$$\therefore \angle ADE = \angle ABC$$

$$\Rightarrow DE \parallel BC$$



Question 18:


ΔABC is an isosceles triangle in which $AB = AC$. D and E are the mid points of AB and AC respectively.



$\therefore DE \parallel BC$
 $\Rightarrow \angle ADE = \angle ABC$ (i)
 Also, $AB = AC$ [Given]
 $\Rightarrow \angle ABC = \angle ACB$ (ii)
 $\therefore \angle ADE = \angle ACB$ [From (i) and (ii)]
 Now, $\angle ADE + \angle EDB = 180^\circ$ [ADB is a straight line.]
 $\therefore \angle ACB + \angle EDB = 180^\circ$
 \Rightarrow The opposite angles are supplementary.
 $\Rightarrow D, B, C$ and E are concyclic
 i.e. D, B, C and E is a cyclic quadrilateral.


Question 19:

Let $ABCD$ be a cyclic quadrilateral and let O be the centre of the circle passing through A, B, C, D . Then each of AB, BC, CD and DA being a chord of the circle, its right bisector must pass through O .
 \therefore the right bisectors of AB, BC, CD and DA pass through O and are concurrent.



Question 20:

$ABCD$ is a rhombus.
 Let the diagonals AC and BD of the rhombus $ABCD$ intersect at O .
 But, we know, that the diagonals of a rhombus bisect each other at right angles.
 So, $\angle BOC = 90^\circ$
 $\therefore \angle BOC$ lies in a circle.




Thus the circle drawn with BC as diameter will pass through O .

Similarly, all the circles described with AB, AD and CD as diameters will pass through O .

Question 21:

ABCD is a rectangle.

Let O be the point of intersection of the diagonals AC and BD of rectangle ABCD.



Since the diagonals of a rectangle are equal and bisect each other,

$$\therefore OA = OB = OC = OD$$

Thus, O is the centre of the circle through A, B, C, D.


Question 22:

Let A, B, C be the given points.

With B as centre and radius equal to AC draw an arc.

With C as centre and AB as radius draw another arc,

which cuts the previous arc at D.



Then D is the required point BD and CD.

In $\triangle ABC$ and $\triangle DCB$

$$AB = DC$$

$$AC = DB$$

$$BC = CB \quad [\text{common}]$$

$$\therefore \triangle ABC \cong \triangle DCB \quad [\text{by SSS}]$$

$$\Rightarrow \angle BAC = \angle CDB \quad [\text{C.P.C.T}]$$

Thus, BC subtends equal angles, $\angle BAC$ and $\angle CDB$ on the same side of it.

\therefore Points A, B, C, D are concyclic.

Question 23:

ABCD is a cyclic quadrilateral

$$\angle B - \angle D = 60^\circ \quad \dots\dots(i)$$

$$\text{and} \quad \angle B + \angle D = 180^\circ \quad \dots\dots(ii)$$

Adding (i) and (ii) we get,

$$2\angle B = 240^\circ$$

$$\therefore \angle B = \frac{240}{2} = 120^\circ$$

Substituting the value of $\angle B = 120^\circ$ in (i) we get


$$120^\circ - \angle D = 60^\circ$$

$$\Rightarrow \angle D = 120^\circ - 60^\circ = 60^\circ$$

The smaller of the two angles i.e. $\angle D = 60^\circ$

Question 24:

ABCD is a quadrilateral in which $AD = BC$ and $\angle ADC = \angle BCD$
 Draw $DE \perp AB$ and $CF \perp AB$



Now, in $\triangle ADE$ and $\triangle BCF$, we have

$$\begin{aligned} \angle AED &= \angle BFC && \text{[each equal to } 90^\circ\text{]} \\ \angle ADE &= \angle ADC - 90^\circ = \angle BCD - 90^\circ = \angle BCF \\ AD &= BC && \text{[given]} \end{aligned}$$

Thus, by Angle-Angle-Side criterion of congruence, we have:

$$\therefore \triangle ADE \cong \triangle BCF \quad \text{[by AAS congruence]}$$

The corresponding parts of the congruent triangles are equal.

$$\therefore \angle A = \angle B$$

Now, $\angle A + \angle B + \angle C + \angle D = 360^\circ$

$$\Rightarrow 2\angle B + 2\angle D = 360^\circ$$

$$\Rightarrow 2(\angle B + \angle D) = 360^\circ$$

$$\Rightarrow \angle B + \angle D = \frac{360}{2} = 180^\circ$$

\therefore ABCD is a cyclic quadrilateral.


Question 25:

If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.

$$\Rightarrow \angle BAD = \angle DCF = 75^\circ$$

$$\therefore \angle DCF = x = 75^\circ$$

$$\therefore x = 75^\circ$$



The opposite angles of the opposite angles of a cyclic quadrilateral is 180°

$$\Rightarrow \angle DCF + \angle DEF = 180^\circ$$

$$\Rightarrow 75^\circ + \angle DEF = 180^\circ$$

$$\Rightarrow \angle DEF = 180^\circ - 75^\circ = 105^\circ$$

$$\text{As } \angle DEF = y^\circ = 105^\circ$$

$$\therefore x = 75^\circ \text{ and } y = 105^\circ$$

Question 26:

Given: Let ABCD be a cyclic quadrilateral whose diagonals AC and BD intersect at O at right angles.

Let $OL \perp AB$ such that LO produced meets CD at M.



To Prove: $CM = MD$

$$\text{Proof: } \angle 1 = \angle 2 \quad \text{[angles in the same segment]}$$

$$\angle 2 + \angle 3 = 90^\circ \quad \text{[}\because \angle OLB = 90^\circ\text{]}$$

$$\angle 3 + \angle 4 = 90^\circ \quad \left[\begin{array}{l} \because \text{LOM is a straight line} \\ \text{and } \angle BOC = 90^\circ \end{array} \right]$$

$$\therefore \angle 2 + \angle 3 = \angle 3 + \angle 4$$

$$\Rightarrow \angle 2 = \angle 4$$

$$\text{Thus, } \angle 1 = \angle 2$$

$$\text{and } \angle 2 = \angle 4$$

$$\Rightarrow \angle 1 = \angle 4$$


$$\therefore OM = CM$$

$$\text{Similarly, } OM = MD$$

$$\text{Hence, } CM = MD.$$

Question 27:


Chord AB of a circle is produced to E.
 If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.
 $\therefore \text{Ext. } \angle BDE = \angle BAC = \angle EAC \dots (1)$



Chord CD of a circle is produced to E.
 $\therefore \text{Ext. } \angle DBE = \angle ACD = \angle ACE \dots (2)$
 Consider the triangles $\triangle EDB$ and $\triangle EAC$.
 $\angle BDE = \angle CAE$ [from(1)]
 $\angle DBE = \angle ACE$ [from(2)]
 $\angle E = \angle E$ [common]
 $\therefore \triangle EDB \sim \triangle EAC$.

Question 28:


Given: AB and CD are two parallel chords of a circle BDE and ACE are straight lines which intersect at E.
 If one side of a cyclic quadrilateral is produced then the exterior angle is equal to the interior opposite angle.
 $\therefore \text{Ext. } \angle ZEDC = \angle A$ and $\text{Ext. } \angle ZDCE = \angle B$



Also, $AB \parallel CD$
 $\Rightarrow \angle ZEDC = \angle B$
 and $\angle ZDCE = \angle A$
 $\therefore \angle A = \angle B$
 $\therefore \triangle AEB$ is isosceles.

Question 29:

AB is a diameter of a circle with centre O. ADE and CBE are straight lines, meeting at E, such that $\angle BAD = 35^\circ$ and $\angle BED = 25^\circ$.
 Join BD and AC.



(i) Now, $\angle BDA = 90^\circ = \angle EDB$ [angle in a semi circle]
 $\Rightarrow \angle EBD = 180^\circ - (\angle EDB + \angle BED)$
 $= 180^\circ - (90^\circ + 25^\circ)$
 $= 180^\circ - 115^\circ = 65^\circ$
 $\therefore \angle DBC = (180^\circ - \angle EBD)$
 $= 180^\circ - 65^\circ = 115^\circ$
 $\therefore \angle DBC = 115^\circ$
 (ii) Again, $\angle DCB = \angle BAD$ [angle in the same segment]
 Since, $\angle BAD = 35^\circ$
 $\therefore \angle DCB = 35^\circ$
 (iii) $\angle BDC = 180^\circ - (\angle DBC + \angle DCB)$
 $= 180^\circ - (\angle DBC + \angle BAD)$
 $= 180^\circ - (115^\circ + 35^\circ)$
 $= 180^\circ - 150^\circ = 30^\circ$
 $\therefore \angle BDC = 30^\circ$